

Introduction of PROSTROLANE B series

Prostrolane B series is new innovative injectables enhanced by Novel Peptide Technology and Sustained Release Technology for Anti Aging, and Body Sliming and Skin Brightening




- *Fat reduction for body slimming*

Inner-B

Let your inner beauty take a shape with
Prostrolane Inner B




Sustained Release Technology



Let your inner beauty take a shape with **Prostrolane Inner B**

Sustained Release Technology



Prostrolane Inner B injectable gel is indicated for deep dermis implantation for lipolysis of moderate to severe double chin, abdomen, buttocks, back of the thighs. Prostrolane Inner B lipolysis injection aims for reduction of localized fat accumulations.

Active Ingredients

Nonapeptide-32, Pentapeptide-43, Tripeptide-41, Octapeptide-11

Capacity

2ml gel in prefilled syringe

Recommended dosage and sessions : 1 session every 2 weeks

Total session : 4 sessions

Advantages

- Reduction of localized fat accumulations - Lipolysis - Inhibition of Lipogenesis

Injection Area: Double Chin area

Injection Volume Double Chin: 4ml per session (Maximum 5ml)

Protocol: 4 sessions at 2 weeks intervals

- Prepare syringe with Lipolysis Injection
- Clean the target area with alcohol swab and well dry the area.
- (If necessary), mark the injection point with erasable marker.
- Inject 0.2ml (cc) to every targeted point. Inject slightly askew angle.



Inner-B



Double Chin

Abdomen

Back of Thigh

Injection Depth

subcutaneous layer of targeted area

Injection Volume

a. Abdomen: 12ml per session (Maximum 20ml)

b. Double Chin: 4ml per session (Maximum 5ml)

Inject 0.2ml (cc) to every targeted point

Recommendation

- Recommend 1~2 times per week Multi-polar RF or Ultra Sound machine to massage the injected area for around 20~30minutes.
- Recommend to do Treadmill Work out for 20~30minutes everyday


Inner-B

Fat Reduction for Body Sliming

2 • Inhibition of Lipogenesis

- Inhibition of Lipid Accumulation

- Cell line: 3T3-L1 pre-adipocyte cell line
- Method: Oil red O staining
- Sample: 0.1-1 $\mu\text{g/ml}$ Inner B complex
- 100ng/ml TNF- α (positive control)
- Culture period: 6days






Inner B complex decreased lipid droplet formation which implied anti-differentiation effect on pre-adipocytes.

Inner-B

Fat Reduction for Body Sliming

3 • Reduction of Localized fat accumulation

Representative micro-CT scanning images





Visceral Fat(mm²) : 4265.3 / Subcutaneous Fat(mm²) : 1258.0

Visceral Fat(mm²) : 5088.1 / Subcutaneous Fat(mm²) : 1465.8

PROSTROLANE *Inner B*

*Innovative Dermal Filler
with Sustained Release Technology & Novel Peptide Technology
for Anti-Obesity*



Clinical Study Before & After

Double chin

2 syringes per session(3 sessions)



Double chin

2 syringes per session(3 sessions)



Double chin

2 syringes per session(3 sessions)



Double chin

2 syringes per session(3 sessions)



Double chin

2 syringes per session(3 sessions)




Double Chin

2 syringes per session (3 sessions)



Double Chin

2 syringes per session (4 sessions)



Double chin

2 syringes per session (4 sessions)



Double chin

2 syringes per session (3 sessions)



Abdomen

4 syringes per session (3 sessions)
Result: -3 cm




Abdomen

4 syringes per session (3 sessions)



Abdomen

4 syringes per session (3 sessions)



Abdomen

4 syringes per session (4 sessions)

Before



After 3 session



After 4 session



Abdomen

4-5 syringes per session (3 sessions)



Flank

2 syringes per session (3 sessions)



Upper Arms

2 syringes per session (3 sessions)



Thigh

4 syringes per session (3 sessions)





Double chin

2 syringes per session(3 sessions)

Prostolane Inner B Double Chin Protocol: 3 sessions, 2 syringes/session

BEFORE



AFTER




Double chin

2 syringes per session(3 sessions)

Prostolane Inner B Double Chin Protocol: 3 sessions, 2 syringes/session

BEFORE



AFTER



Abdomen

4-5 syringes per session (3 sessions)

Protocol: 6 syringes/session; 3 sessions

Body measurements: Waist 81,5 cm/ Umbilicus 97 cm/ Hips 104,5 cm

Before



15 days



30 days



Final measurements: Waist 80 cm/ Umbilicus 91 cm/ Hips 101 cm

